

On Classical Ideal Gases

Jacques ARNAUD ^{*}, Laurent CHUSSEAU [†], Fabrice PHILIPPE [‡]

June 19, 2012

Abstract

We show that the thermodynamics of ideal gases may be derived solely from the democritian concept of corpuscles moving in vacuum plus a principle of simplicity, namely that these laws are *independent* of the laws of motion, aside from the law of energy conservation. Only a single corpuscle in contact with a heat bath submitted to a z and t -invariant force is considered. Most of the end results are known but the method appears to be novel. The mathematics being elementary the present paper should facilitate the understanding of the ideal-gas law and of classical thermodynamics even though not-usually-taught concepts are being introduced. Appendices clarify detailed points.

^{*}Mas Liron, F30440 Saint Martial, France

[†]Institut d'Électronique du Sud, UMR n°5214 au CNRS, Université Montpellier II, F34095 Montpellier, France

[‡]LIRMM, UMR n°5506 au CNRS, 161 rue Ada, F34392 Montpellier, France

1 Introduction

Let us first quote Feynman [1]: “If, in some cataclysm, all of scientific knowledge were to be destroyed, and only one sentence passed to the next generations of creatures, what statement would contain the most information in the fewest words? I believe it is the hypothesis that all things are made of atoms. In that one sentence there is an enormous amount of information about the world, if just a little imagination and thinking are applied”. In line with this quotation, it seems worthwhile exploring how much physics may be derived from the corpuscular concept. Only some qualitative observations are needed. One is the fact that it takes some time, denoted here $\tau(z_m)$, for a corpuscle thrown upward on earth to reach an altitude z_m and come back to the ground level. We have of course $\tau(0) = 0$. If the corpuscle bounces elastically on the ground, $\tau(z_m)$ represents the oscillation period. The time during which the corpuscle is located above some altitude $z \leq z_m$ during a period is: $\tau(z_m - z)$. Under our assumptions the τ -function does not depend on the initial altitude or initial time. A second observation is that two bodies left in contact for a sufficient period of time tend to reach the same temperature, as one can judge by our senses.

The purpose of the present paper is thus to show that the thermodynamics of ideal gases and in particular the ideal-gas law may be obtained on the sole basis of the democritian model according to which nature consists of corpuscles moving in a vacuum, plus a principle of simplicity: namely that these fundamental laws are *independent* of the law of corpuscle motion (non-relativistic, special relativistic [2], or otherwise). The temperature θ enters solely for dimensional reasons. We later show from the general expressions of the gas internal energy and of the force (or pressure) that the heat delivered by the gas is θdS , an expression of the entropy S being given. This result enables us to prove that the formally-introduced temperature θ is a thermodynamic temperature. Indeed, we recover for ideal gases the general Carnot result asserting that the efficiency of reversible thermal engines is: $1 - \theta_l/\theta_h$, where θ_l denotes the cold bath temperature and θ_h the hot bath temperature. Since Kelvin time this expression *defines* the thermodynamics temperature to within an arbitrary proportionality factor, which is fixed by specifying that $\theta \approx 370 \times 10^{-23}$ joules at the water triple point. An alternative derivation of the Carnot result, also solely based on the concept of potential energy, is in [3, 4].

In the title of this paper the word “classical” means: “non-quantum” ($\hbar \rightarrow 0$), but the non-relativistic approximation is not made, except in examples. “Ideal Gases” refer to non-interacting identical corpuscles. Accordingly, the corpuscles are independent, and we may consider a single corpuscle. As a result of the slight quivering of the container wall, there is an exchange of energy between the corpuscle and the heat bath (for a static wall, even a rough one, the corpuscle energy would be a constant of motion). Thus we are working in the so-called “canonical ensemble” [5]. The paper [6]: “Microcanonical single-particle distributions for an ideal gas in a gravitational field” is valuable because it establishes (within the non-relativistic approximation) the gas law in the micro-canonical ensemble with the effect of gravity taken into account as in the present paper. These authors find that the micro-canonical result approaches the canonical result for few corpuscles only. However that paper is complicated both from the mathematical and physical viewpoints. The reader may wish to compare the micro-canonical many-corpuscle paper just cited with the present canonical single-corpuscle paper in terms of pedagogical value. Of course the final results are the same.

The reader may feel that our statement that the above invariance principle *implies* the ideal-gas laws, without anything else, is surprising. Yet, we hope that we can convince him/her that this is indeed the case. On the other hand, a quick reading of standard books on Thermodynamics may

lead other readers to believe that this is a well-known fact. For example, Callen [7] states correctly that: “The essence of the ideal-gas law is that molecules of the gas do not interact. This simple fact *implies* that $PV \propto NT$ ”. However, in order to reach this conclusion, the author needs postulate the Newtonian law of motion, quantum theory, and the Boltzmann factor. The claim that corpuscle independence entails the ideal-gas law (without anything else) therefore does not appear to have been justified before.

Let us emphasize that our goal is to derive the ideal-gas laws from first principles, only conservation of potential energy and thermal equilibrium being assumed (our model cannot describe directly the so-called “adiabatic atmospheres” that are in mechanical but not thermal equilibrium). We do not use the concept of kinetic energy, nor do we postulate any particular law of corpuscle motion. Accordingly, a given potential $\phi(z) = wz$ for a weight w does not imply any specific law of motion $z(t)$. Because the presence of a corpuscle affects negligibly the potential one may consider the latter to be an *external* potential. To our knowledge, the concepts presented in this paper have not been considered before. However, the final expressions relating to the relativistic ideal-gas law in the presence of gravity are known, see for example [8].

Admittedly the present paper departs much from usually-taught concepts in Thermodynamics [9]. We indeed ignore the concepts of kinetic energy, of micro-states, and the laws of Physics that would provide the motion $z = z(t)$ (to within arbitrary t and z translations) of corpuscles of given energy. The temperature θ (to within a constant factor) follows from dimensional considerations and not as usual from the derivative of the average energy $U(S, h)$ with respect to the entropy S at constant volume (here denoted h). Quite to the contrary, an explicit expression of the (classical ideal-gas) entropy function $S(\theta, h)$ is derived from the average force $\langle F \rangle$ (or pressure) and average energy U . This change of perspective suggests that one should first emphasize, in teaching the subject to students, the Democritian concept of corpuscles moving in vacuum and the concept of potential energy (e.g., from cords and pulleys experiments). Next, consider the up and down motion of a corpuscle thrown upward from $z = 0$ in the earth gravitational field (supposed to be static and independent of the vertical z coordinate), the particular function $z(t)$ being left undefined. As a result of weak thermal contact with the bath (that needs not be specified further), the maximum altitude z_m varies slowly in the course of time. This variation is described statistically by some probability law $\omega(z_m)$. It should be explained next that a weight measured by a high-inertia balance does not depend on possible up and down motions of the weight. Accordingly, for a unit weight, the weight impact is equal to the motion period. This observation enables us to express the force F in terms of round-trip times when a rigid plate is present at $z = h$, see Fig 1. The condition that the average force be *independent* of the $\tau(z_m)$ -function introduced above provides the ideal-gas law in full generality.

To conclude, the approach advocated in this paper enables us to recover classical results in Thermodynamics in a considerably generalized form (but only for independent classical corpuscles and laws of Physics that are independent of time and altitude translations). Admittedly, our presentation goes against the order in which basic concepts are usually taught. But the greater simplicity and generality will perhaps be appreciated.

2 Average force exerted by a corpuscle on a piston

We consider a unit-area cylinder with vertical z -axis in uniform gravity resting on the ground ($z = 0$) at some temperature. A tight piston is free to move in the vertical direction. The cylinder height is denoted by h and contains a single corpuscle of weight $w = 1$. In our one-dimensional




Figure 1: Space-time (z, t) trajectory of a corpuscle of unit weight ($w = 1$) bouncing off the ground ($z = 0$). The maximum altitude reached by the corpuscle is $z_m = E$, where E denotes the energy. The motion is periodic with period $\tau(z_m)$, where $\tau(Z)$ denotes the corpuscle round-trip time at a distance Z from the top of the trajectory. When the altitude is restricted to h by a plate (dashed horizontal line) the motion remains periodic with a period evidently equal to: $\tau(z_m) - \tau(z_m - h)$. Note that this expression holds even if the motion is not symmetric in time.

model, the pressure P corresponds to the average force $\langle F \rangle$, the volume V to the height h , and $N = 1$. Our result provides the ideal-gas law in a generalized form, taking into account gravity. In that case, the pressure varies as a function of altitude. More precisely, the force exerted by the corpuscle on the lower end of the cylinder exceeds the force exerted on the upper end (or piston) by the corpuscle weight. But in the absence of gravity, the forces exerted on both ends would be the same. We are introducing (static and uniform) gravity mainly because this helps clarify the concept of corpuscle energy: the corpuscle energy E is the maximum altitude z_m that the corpuscle would reach in the absence of the piston when the corpuscle weight is unity. The corpuscle bounces elastically off the ground and off the piston (if it reaches it) that is, without any loss or gain of energy.

Consider first the case where h is infinite, that is, in the absence of a piston. The time period is denoted as before by $\tau(Z)$ with $Z = z_m$. The average force exerted on the ground, equal to the corpuscle weight $w = 1$, is the impulse i divided by the period. Thus $1 = i/\tau(z_m)$ or $i = \tau(z_m)$. In other words, the impulse transmitted to a piston when the corpuscle impacts on it is equal to the motion period. If the piston is located at $z = h$ the impulse it receives is: $i_h = \tau(z_m - h)$ if $z_m \geq h$, and zero otherwise. On the other hand, the motion period is $\tau(z_m)$ if $z_m \leq h$ and $\tau(z_m) - \tau(z_m - h)$ if $z_m > h$.

Accordingly, the force F experienced by the piston when the corpuscle energy is z_m is the ratio of the impulse and period:

$$\begin{cases} F(z_m) = 0 & z_m \leq h, \\ F(z_m) = \frac{i_h}{\tau(z_m) - \tau(z_m - h)} = \frac{\tau(z_m - h)}{\tau(z_m) - \tau(z_m - h)} & z_m > h. \end{cases} \quad (1)$$

Because the cylinder lower end is in contact with a bath, there is a slight quivering (thermal motion) and the corpuscle energy z_m slowly varies in the course of time. Accordingly, the average force $\langle F \rangle$ experienced by the piston is, if $\omega(z_m)$ denote the energy distribution:

$$\langle F \rangle = \frac{\int_0^\infty dz_m \omega(z_m) F(z_m)}{\int_0^\infty dz_m \omega(z_m)} = \frac{\int_h^\infty dz_m \omega(z_m) \frac{\tau(z_m - h)}{\tau(z_m) - \tau(z_m - h)}}{\int_0^h dz_m \omega(z_m) + \int_h^\infty dz_m \omega(z_m)}. \quad (2)$$

According to our simplicity principle, the average force $\langle F \rangle$ must be *independent* of the corpuscle equation of motion, and thus of the $\tau(\cdot)$ -function. This condition obtains from (2) if one selects the following energy distribution:

$$\begin{cases} \omega(z_m) = \exp(-z_m/\theta) \tau(z_m) & z_m \leq h, \\ \omega(z_m) = \exp(-z_m/\theta) (\tau(z_m) - \tau(z_m - h)) & z_m > h, \end{cases} \quad (3)$$

where θ has the dimension of an energy when non-unity weights w are considered. The average force becomes, using (2) and (3):

$$\begin{aligned} \langle F \rangle &= \frac{\int_h^\infty dz_m \exp(-z_m/\theta) \tau(z_m - h)}{\int_0^h dz_m \exp(-z_m/\theta) \tau(z_m) + \int_h^\infty dz_m \exp(-z_m/\theta) (\tau(z_m) - \tau(z_m - h))} \\ &= \frac{w}{\exp(w h/\theta) - 1} \end{aligned} \quad (4)$$

if we restore w (previously set equal to unity). In the above integrals going from h to ∞ we have replaced $\exp(-z_m/\theta)$ by $\exp(-h/\theta) \exp(-(z_m - h)/\theta)$ and introduced the variable $z'_m \equiv z_m - h$, so that all the integrals go from zero to infinity and cancel out. Note that even though integral signs have been introduced no integration has been performed. We simply consider an integral as a sum of terms and employ the rule of addition associativity. We have employed also the fact that $f(x) = \exp(x)$ is the only function such that $f(a + b) = f(a)f(b)$.

For a collection of N independent corpuscles having weights w_i , $i = 1, \dots, N$ respectively, the force is a sum of N terms of the form given in (4). In the case of zero gravity ($w=0$ or more precisely: $wh \ll \theta$), the above expression gives: $\langle F \rangle h = \theta$. Thus we have obtained the ideal-gas law: $\langle F \rangle h = N \theta$.

Average force for a three-dimensional space: We suppose that the cylinder radius is very large compared with h , and we do not consider the force exerted by the corpuscle on the cylinder wall. Motion of the corpuscle along directions perpendicular to z (say, x and y) *does* affect the round-trip time function $\tau(Z)$. However, since the average force does not depend on this function, the ideal-gas law is unaffected. This is so for any physical system involving a single corpuscle provided the physical laws are invariant under a z -translation (besides being static).

The internal energy, to be discussed in the following section, though, is incremented. One can prove that in the non-relativistic approximation and in the absence of gravity the internal energy is multiplied by 3. It would be incremented further by corpuscle rotation or vibration, not considered here. Using conventional methods, Landsberg [8] in Eq. (2.6), and Louis-Martinez [10] in Eq. (68), obtain exactly the same result as given above (except for the factor 3 in the expression of the internal energy, relating to the number of space dimensions considered).

3 Internal energy

The gas internal energy U is the average value of E , the gravitational energy being accounted for. Note that only corpuscule motion along the z -axis is being considered. In the present section a non-unity (but constant) weight w is considered to clarify the dimensions of the quantities introduced. The expression of U is, using the energy distribution given in (3):

$$\begin{aligned}
U &= \frac{\int_0^{wh} dE E \exp(-E/\theta) \tau(E/w) + \int_{wh}^{\infty} dE E \exp(-E/\theta) (\tau(E/w) - \tau(E/w - h))}{\int_0^{wh} dE \exp(-E/\theta) \tau(E/w) + \int_{wh}^{\infty} dE \exp(-E/\theta) (\tau(E/w) - \tau(E/w - h))} \\
&= \frac{\int_0^{\infty} dE E \exp(-E/\theta) \tau(E/w) - \int_{wh}^{\infty} dE (E - wh + wh) \exp(-E/\theta) \tau(E/w - h)}{\int_0^{\infty} dE \exp(-E/\theta) \tau(E/w) - \int_{wh}^{\infty} dE \exp(-E/\theta) \tau(E/w - h)} \\
&= \frac{(1 - \exp(-wh/\theta)) \int_0^{\infty} dE E \exp(-E/\theta) \tau(E/w) - wh \int_{wh}^{\infty} dE \exp(-E/\theta) \tau(E/w - h)}{(1 - \exp(-wh/\theta)) \int_0^{\infty} dE \exp(-E/\theta) \tau(E/w)} \\
&= \frac{(1 - \exp(-wh/\theta)) \int_0^{\infty} dE E \exp(-E/\theta) \tau(E/w) - wh \exp(-wh/\theta) \int_0^{\infty} dE \exp(-E/\theta) \tau(E/w)}{(1 - \exp(-wh/\theta)) \int_0^{\infty} dE \exp(-E/\theta) \tau(E/w)} \\
&= \frac{\int_0^{\infty} dE E \exp(-E/\theta) \tau(E/w)}{\int_0^{\infty} dE \exp(-E/\theta) \tau(E/w)} - \frac{wh}{\exp(wh/\theta) - 1}. \tag{5}
\end{aligned}$$

The internal energy divided by the temperature: $U(\theta, h)/\theta$, thus is the sum of a term function of θ but not of h , and a term which tends to $-\theta$ when $wh \ll \theta$. To evaluate the first term we need to know the round-trip time $\tau(Z)$ to within an arbitrary proportionality factor, and an integration must be performed in that case.

In order to write the above expression in terms of dimensionless quantities, we must introduce a universal constant, c , having the dimension of speed. Defining: $x \equiv E/\theta$, $\theta' \equiv \theta/mc^2$, $h' \equiv h/(c^2/g)$, (5) becomes, remembering that $w = mg$:

$$\frac{U}{\theta} = \frac{\int_0^{\infty} dx x \exp(-x) f(\theta' x)}{\int_0^{\infty} dx \exp(-x) f(\theta' x)} - \frac{h'/\theta'}{\exp(h'/\theta') - 1}, \tag{6}$$

where the function $f(y)$ depends on the round-trip time.

In the special case of non-relativistic motion $y \ll 1$, we have $f(y) \propto \sqrt{y}$. The first term in (6) is then [11] equal to $3/2$, so that in the absence of gravity the internal energy U is equal to $3\theta/2 - \theta = \theta/2$, a well-known result. More generally we have: $f(y) \propto \sqrt{y^2 + 2y}$. A plot of U from low ($\theta \ll mc^2$) to high ($\theta \sim mc^2$) temperatures can be seen in [4]. A three-dimensional expression of U in terms of Bessel functions can be found, for example, in Greiner, p. 234. [12].

4 The energy θ is a thermodynamic temperature

We prove in this section that θ , introduced in previous sections on dimensional grounds only, is a thermodynamic temperature. We do this by showing that the efficiency of a reversible thermal cycle employing ideal gases is: $1 - \theta_l/\theta_h$, where θ_l is the cold-bath temperature and θ_h the hot bath temperature: this is the accepted Kelvin definition of absolute temperatures.

The expressions given earlier for the average force $\langle F \rangle$ in (4) and the internal energy U in (5) may be written, setting $\beta \equiv 1/\theta$, as:

$$\begin{aligned}\langle F \rangle &= \frac{\partial \ln(Z)}{\beta \partial h} & U &= -\frac{\partial \ln(Z)}{\partial \beta} \\ Z(\beta, h) &= (\exp(-\beta w h) - 1) \int_0^\infty dE \exp(-\beta E) \tau(E/w).\end{aligned}\tag{7}$$

Z is essentially the quantity called in statistical mechanics the partition function. It becomes dimensionless if it is divided by the reduced Planck constant $\hbar$, which however plays here no physical role.

If we introduce the Helmholtz free-energy (the letter A is from the German “Arbeit” or work): $A(\theta, h) \equiv -\theta \ln(Z(\theta, h))$ the expressions in (7) are conveniently written:

$$\langle F \rangle = -\frac{\partial A}{\partial h} \quad U = A - \theta \frac{\partial A}{\partial \theta}.\tag{8}$$

From (8) we obtain:

$$-\delta Q \equiv dU + \langle F \rangle dh = dA - \frac{\partial A}{\partial \theta} d\theta - \frac{\partial A}{\partial h} dh - \theta d\left(\frac{\partial A}{\partial \theta}\right) = \theta dS \quad S \equiv -\frac{\partial A}{\partial \theta},\tag{9}$$

where δQ represents the heat released by the gas, from the law of conservation of energy. For any function $f(\theta, h)$ such as U , A , S : $df \equiv \frac{\partial f}{\partial \theta} d\theta + \frac{\partial f}{\partial h} dh$. Note that we employ only two independent variables, namely θ and h , so-that partial derivatives are un-ambiguous. If the gas is in contact with a thermal bath ($\theta=\text{constant}$), δQ is the heat gained by the bath. The quantity S defined above is called “entropy”. In particular, if heat cannot go through the gas container wall (adiabatic transformation) we have $\delta Q = 0$ that is, according to the above result: $dS = 0$. Thus reversible adiabatic transformations are isentropic.

The Carnot cycle: A Carnot cycle consists of two isothermal transformations at temperatures θ_l and θ_h , and two intermediate reversible adiabatic transformations ($dS = 0$). After a complete cycle the entropy recovers its original value and therefore $dS_l + dS_h = 0$. According to (9): $-\delta Q_l = \theta_l dS_l$, $-\delta Q_h = \theta_h dS_h$ and therefore $\delta Q_l/\theta_l + \delta Q_h/\theta_h = 0$. Energy conservation gives the work δW performed over a cycle from: $\delta W + \delta Q_l + \delta Q_h = 0$. The cycle efficiency is defined as the ratio of δW and the heating $-\delta Q_h$ supplied by the hot bath. We have therefore: $\eta \equiv \frac{\delta W}{-\delta Q_h} = \frac{\delta Q_h + \delta Q_l}{-\delta Q_h} = 1 - \frac{\theta_l}{\theta_h}$, from which we conclude that θ is the “thermodynamic temperature”. Since Kelvin time, thermodynamics temperatures are strictly defined from Carnot (or other reversible) cycles efficiency. In practice, temperatures may be measured by other means and employed in other circumstances.

We have implicitly assumed in the above discussion that the working medium (presently an ideal gas) has reached the bath temperature before being contacted with it. Otherwise, there would be at that time a jump in entropy, and the cycle would no longer be reversible. Given initial θ , h values, the temperature change $d\theta$ for an increment dh in the isentropic regime ($dS = 0$) follows from the relation: $d\theta = -\left(\frac{\partial S/\partial h}{\partial S/\partial \theta}\right)dh$, where $S(\theta, h)$ may be expressed in terms of $Z(\theta, h)$, (7), from the above expressions. The details will be omitted. It suffices to know that θ may be varied by varying h , in a calculable manner, in an isentropic transformation.

Practical units: The energy $\theta = \langle F \rangle h$ has been defined so far only to within a multiplicative factor from dimensional considerations. This factor is fixed by agreeing that $\theta = 273.16 k_B$ exactly when the cylinder is in thermal equilibrium with water at its triple point. Here $k_B = 1.38066... 10^{-23}$ joules, is considered as an energy unit (akin to the calorie = 4.182... joules). This manner of defining θ is equivalent to the usual one, though expressed differently. The dimensionless quantity $T \equiv \theta/k_B$ is the usual unit of thermodynamic temperature, expressed in kelvin.

Next, measurements have shown that the number of atoms in 0.012 kg of carbon 12 is: $N_A \approx 6.0221367 10^{23}$. For this quantity of matter (called a mole), the ideal-gas law therefore reads: $\langle F \rangle h = N_A \theta$, or: $PV = RT$, with the ideal-gas constant: $R \equiv N_A k_B \approx 8.31451$ joules per kelvin per mole.

5 Conclusion

Let us briefly recall the concepts introduced in the present paper. One can imagine that after having introduced the corpuscular concept, Democritus observed the elastic bounces of a unit weight on a balance and defined the weight “impulse” from the motion period. Not knowing the nature of the motion (parabolic? hyperbolic?), he may have thought of introducing a weight factor such that the average force $\langle F \rangle$ *does not depend* on the law of motion. This, as we have seen, may be done. This weight factor involves for dimensional reasons a quantity θ having the dimension of energy. Considering a thermal engine operating between two baths at temperatures θ_l, θ_h one finds on the basis of the principles just stated that the maximum efficiency is: $1 - \theta_l/\theta_h$. This allows us to call θ the thermodynamic temperature.

The present paper provides a first-principle proof of the ideal-gas law, including a possible effect of uniform gravity with no knowledge of the round-trip time function $\tau(Z)$ being required. Our thesis is that this law may be obtained on the sole basis of the Democritus model of corpuscles and vacuum. It is indeed unnecessary to specify the law $z(t; E)$ of corpuscle motion, where E denotes the energy. Explicit expressions of the internal energy are obtained provided the round-trip time be known. One can further show that the ideal-gas internal energy depends only on temperature in the absence of gravity. The theory presented is strictly classical. As such, it does not depend on the numerical value of the universal constant $\hbar$. However, the introduction of this quantity is needed to make the results dimensionless.

A History and early experiments

In this appendix we recall some early concepts relating to heat, fundamental experimental results reported around the 17th century, and traditional ways of describing these results.

Concepts relating to heat since the antiquity: From Boyle’s time (*circa* 1660) to Carnot’s time (*circa* 1824), the most popular theory, usually referred to as the “caloric theory of heat”, considered heat as a kind of fluid consisting of weightless mutually repelling particles, attracted by matter. This fluid could flow from hotter to colder matter, see Levermore [13]. This hypothesis, however, did not explain Rumford’s experiments showing that friction alone generates heat. Doubts about the validity of the caloric theory were expressed in Carnot’s notes, published after his early death. As a matter of fact, the alternative model of heat as a form of *matter* motion (without any additional fluid) had already been enunciated in the antiquity.

Democritus, who lived about 300 years B.C., described nature as a collection of corpuscles that cannot be split, moving in vacuum. These corpuscles differ from one-another in form, position and weight. In the case of a gas, interaction between corpuscles may often be neglected, but they collide with the container's walls. Platon [14] ascribed heat to corpuscular motion: "Heat and fire are generated by impact and friction, but that's motion". Much later, Francis Bacon (1561-1626) wrote: "The very nature of heat is motion, and nothing else". This view-point is more explicit in Daniel Bernoulli writing (1738): "Gas atoms are moving randomly, and pressure is nothing else but the impact of the atoms on their container walls". Lastly, Carnot introduced energy considerations *circa* 1830: "Heat is nothing but motive power, or rather another form of motion. When motive power is destroyed, heat is generated precisely in proportion of the motive power destroyed. Likewise, when heat is destroyed, motive power is generated".

Experimental results relating to air The first accurate experiments relating to gases relied on the invention of the thermometer by Galileo and of the barometer by his assistant Torricelli. Then, to Pascal experiments on atmospheric pressure. Pascal ascribed the diminution of the height of a mercury column as a function of altitude to the reduction of the weight of the air above the barometer. It was later shown that the pressure decays exponentially (or more precisely as the sum of two exponentials, one relating to nitrogen, and one relating to oxygen).

Let us recall the crucial experiments performed in the seventeenth century concerning the properties of air. Air, consisting mostly of di-atomic nitrogen, may be viewed as an ideal gas. When a tight box contains some amount of air, the volume-pressure product is a constant at room temperature, a law enunciated for the first time by Boyle in 1660: "Pressure and expansion are in reciprocal proportions". Boyle employed a J-shaped glass tube, with the sealed small side full of air, and the other full of mercury. The J left-side height was a measure of volume and the right-side height a measure of pressure. Subsequent experiments have shown that this law is applicable at any constant temperature, for example at various liquid boiling temperatures, within some experimental range. We call "generalized Boyle law" the expression: $V(P, \theta) = f(\theta)/P$, where P represents pressure, V volume, and $f(\theta)$ some temperature measure.

From an experimental standpoint we could define temperature as the pressure relating to some given amount of matter contained within some fixed volume. As the temperature gets higher the pressure increases. This pressure may be used to define θ . Of course, different temperature scales would be obtained for different substances, but such thermometers may be calibrated one against another because temperatures tend to equalize in equilibrium. Rarefied helium may be described with great accuracy as a collection of independent corpuscles, except perhaps at very low temperatures when quantum effects become significant and at very high temperatures when the helium atoms may get ionized. The theory presented in this paper shows on the basis of the corpuscular model that θ , initially introduced formally from dimensional considerations, coincides with the thermodynamic temperature. This is the temperature that enters in the expression of thermal-engine efficiencies.

Gay-Lussac has shown in 1802 that, at atmospheric pressure, the volume increment of various gases from freezing to boiling water temperatures is 37.5 %. The empirical Gay-Lussac law makes sense only if one specifies which thermometer is being employed. One may employ a gas thermometer from a selected gas such as helium in two ways. One method consists of defining the temperature as the cylinder height (or volume) at a fixed pressure, for example at the standard atmospheric pressure. In the following, we assume that a second method is being employed instead: the temperature is defined as the force that must be exerted on the piston to maintain the height at

a fixed value, for example one meter. If the Gay-Lussac experiment were applied to a gas identical to the gas employed in the thermometer (helium in our example) the fact that pressure is proportional to temperature would be obvious. The importance of the Gay-Lussac experiment is that the proportionality law is found to be approximately valid for any gas. It was subsequently discovered that the Gay-Lussac law is reasonably accurate only at very small pressures. The theoretical reason that explains this observation is that, at low pressures, the molecules of the tested gas and those of the thermometer gas may both be considered as independent non-interacting corpuscles.

In 1803, Dalton, on the basis of his studies of chemical compounds and gaseous mixtures suggested that matter consists of atoms of different masses that combine in simple ratios. He discovered the partial-pressure law according to which the total pressure exerted by a gas mixture is equal to the sum of the pressure that each one of the gases would exert if it occupied the full volume alone. Finally, in 1811, Avogadro concluded that equal volumes of gases at the same temperature and pressure contain the same number of molecules (or corpuscles). This entails that PV/θ is proportional to N , now interpreted as the number of corpuscles. On empirical grounds, the ideal-gas law may therefore be written as:

$$PV = N\theta \tag{10}$$

where V denotes the volume, P the pressure, N the number of corpuscles, and $\theta \equiv k_B T$ the temperature.

Usual kinetic and statistical theories The ideal-gas law has been partly explained on the basis of a kinetic theory by Waterston [15] in 1843, the kinetic theory being based itself on non-relativistic mechanics. Let us now recall the basic assumptions on which rest the usual proofs of the barometric and ideal-gas laws. A recent reference by Miller [16] lists the assumptions on which the gas kinetic theory is supposed to rest. Some of them express the democritian hypothesis and are indeed essential. The usefulness of the others [17] may be questioned (these assumptions are neither necessary nor sufficient). As we shall show, none are needed. It suffices that the (perhaps unique) corpuscle be in thermal contact with the ground. Besides, the condition that the average kinetic energy *depends* only on the system temperature (*cf* [17]) is insufficient to establish the ideal-gas law because the temperature remains undefined. In the non-relativistic approximation the appropriate assumption should read: “The average kinetic energy (defined as the difference between the total energy and the potential energy) is *proportional* to the system temperature”. But in general a temperature so defined, say $\tilde{T}$, is *not* the thermodynamic temperature. That is, the maximum engine efficiencies are not $1 - \tilde{T}_l/\tilde{T}_h$.

Let us sketch the most usual derivations of the barometric and ideal-gas laws to remind the readers of the underlying assumptions. Note that the barometric law may be obtained from the ideal-gas law, and conversely, if one postulates that weightless infinitely-thin plates may be introduced or removed at will in the gas at various altitudes. However, this postulate is at best plausible. Norton [18] derived the ideal-gas law from the barometric equation. But the latter involves the Boltzmann factor that requires other physical considerations (see below), while in the present paper this factor comes in naturally, that is, for purely mathematical reasons. The derivation of the Boltzmann factor is usually based on a quantization of the energy and the postulated equi-probability of the micro-states. Let the discrete (non-degenerate) energy levels be denoted by $\varepsilon_1, \varepsilon_2, \dots$. If distinguishable corpuscles are distributed among the energy levels, with n_1 corpuscles in level 1, n_2 corpuscles in level 2, and so on, the number of ways of doing that is inversely proportional to: $n_1!n_2!\dots$. It is postulated that this number reaches its maximum value at equilibrium

under the constraint that $n_1 + n_2 + \dots = N$, the total number of corpuscles, and $n_1 \varepsilon_1 + n_2 \varepsilon_2 + \dots = E$, the total energy. In the limit of large N values, one finds that: $n_i \propto \exp(-\varepsilon_i/\theta)$, for some θ -value that depends on N and E . Even though physicists are now-a-day very familiar with that procedure, it is not so easy to explain it to students. Besides, it rests on a number of assumptions.

The earliest derivation of the ideal-gas law is based on non-relativistic mechanics. For a one-dimensional model, one considers a corpuscle moving back and forth between two plates separated by a distance h , one of them playing the role of a piston. If v denotes the speed of a corpuscle, an impact on a plate delivers to it an impulse $2mv$ where m denotes the corpuscle mass, and this occurs every $2h/v$ time units. It is concluded that the force F exerted on the piston is: $2mv/(2h/v) = mv^2/h$, that is: $Fh = 2K$, where $K = \frac{1}{2}mv^2$ denotes the kinetic energy. It is recognised that there may be a distribution of kinetic energies. *Postulating* that the temperature θ is proportional to the average kinetic energy one obtains for the average force the ideal-gas law: $\langle F \rangle h \propto \theta$. Alternatively, one may quantize the corpuscle wave-function and employ the Boltzmann factor.

The procedure described above has been generalized to relativistic motion ($k_B T \sim mc^2$), where mc^2 denotes the corpuscle rest energy, but the relation of v to p is different, and averaging requires a generalized Boltzmann factor. The same ideal-gas law is valid at any temperature (within the corpuscular model). Our thesis is that the ideal-gas law has simply nothing to do with the law of corpuscle motion, and that it is therefore not surprising that it holds for the Galileo, Einstein, and many other laws of motion. These are the reasons why we feel that the traditional proofs are unsatisfactory. An alternative was offered in the main text, considering in succession the ideal-gas law and the internal energy. We have finally shown that the formally introduced temperature θ is the thermodynamic temperature. Expressions for the round-trip time of a corpuscle from and back to the ground $z=0$ level, denoted by τ , is derived in Appendix B. The case of potentials not proportional to z (z -dependent weights) was treated by the same method in [19], see Appendix C.

B Equations of motion

In the main text we have shown that the barometric and ideal-gas laws are independent of the corpuscle motion, while the energy depends on the round-trip time function $\tau(Z)$ to within an arbitrary scale factor. In some applications, e.g., for adiabatic atmospheres whose treatment requires an expression of the gas energy density, it is important to know that function. In the present appendix we recall that equations of motion in an inertial reference frame are best formulated through the hamiltonian formulation. We are considering the non-relativistic approximation, the motion of corpuscles that are free (aside from gravity), and a model of gravity based on a “rain” of particles. The formulation in the case of an accelerated frame is mentioned [20].

The main-text results are based on the assumption that, if $z = z(t)$ is a particular solution, then *all* the solutions are of the form: $z - z_o = f(t - t_o)$ where z_o , t_o are constants. In other words, the general solution follows from the particular one initially selected through altitudes and time translations. This condition is satisfied if we set:

$$v(t) \equiv \frac{dz(t)}{dt} = H'(p), \quad \frac{dp(t)}{dt} = -w. \quad (11)$$

where $H(p)$ is a nearly arbitrary function and primes denote derivatives with respect to the argument. Integration of the last equation in (11) gives: $p = -w(t - t_o)$ with t_o an arbitrary constant. The equation for v gives: $z(t) - z_o = -\frac{H(-w(t-t_o))}{w}$.

As one recalls from the main text, $Z \equiv z_m - z > 0$ denotes the distance between the maximum altitude and some altitude z . The round-trip time τ denotes the time duration between two successive equal z -values. It follows from the above $z(t)$ expression that for two w -values, say w_1 and w_2 , the following relation holds [21]

$$\tau_2(Z) = \frac{w_1}{w_2} \tau_1\left(\frac{w_2}{w_1} Z\right), \quad (12)$$

Non-relativistic approximation for a constant weight: Expanding to second order: $H(p) = a_o + a_1 p + \frac{1}{2m} p^2$, where we choose to write a_2 as $1/2m$. The expression given above for $z(t)$ can be rearranged to give under that second-order approximation: $z(t) - \tilde{z}_o = -\frac{w}{2m}(t - \tilde{t}_o)^2 \equiv -\frac{g}{2}(t - \tilde{t}_o)^2$, where we have introduced new integration constants $\tilde{z}_o$, $\tilde{t}_o$, and set: $w/m \equiv g$. Galileo, incidentally, observed that g is the same for all corpuscles; however this experimental fact is unimportant in the main part of the present paper that deals with N identical corpuscles. With the initial conditions: $z(0) = 0$ and $t(0) = 0$, the corpuscle motion is described by: $z(t) = -\frac{1}{2}gt^2$. In the present form of $z(t)$, $z_m = 0$ and we have: $Z = -z$. Thus, $t_{2,1} = \pm\sqrt{2Z/g}$ and $\tau(Z) = t_2 - t_1 = 2\sqrt{2Z/g} \propto \sqrt{Z}$ if g is a constant.

Relativistic expression for constant electrical forces: We presently have for a weightless corpuscle with electrical charge e and submitted to a constant electrical field $\mathcal{E}$, setting $w \equiv e\mathcal{E}$, (see Landau [22], p.115, for the relativistic form without gravity in three dimensions):

$$\begin{aligned} H(z, p) &= mc^2 \sqrt{1 + (p/mc)^2} + wz \rightarrow p = -wt, \\ \frac{dz(t)}{dt} &= \frac{\partial H(z, p)}{\partial p} = c \frac{\tilde{p}}{\sqrt{1 + \tilde{p}^2}} \quad \tilde{p} \equiv \frac{p}{mc}. \end{aligned} \quad (13)$$

From these expressions we may obtain the equation of motion with $z(0) = z_m$, and the round-trip time as a function of $z_m - z$. Thus, if we set for brevity $m = 1$, $g = 1$, $c = 1$, we have: $H(p) = \sqrt{1 + p^2}$, $v \equiv dz/dt = p/\sqrt{1 + p^2} = -t/\sqrt{1 + t^2}$, or $z = 1 - \sqrt{1 + t^2}$. It follows that, according to our definitions of τ and Z given in the main text, $\tau(Z) = 2\sqrt{Z^2 + 2Z}$. Introducing this function in the main-text expression one find that the energy may be expressed in terms of Bessel functions, a well-known result [23]. The non-relativistic approximation corresponds to neglecting Z^2 compared with Z in that expression.

Note that corpuscles in free-space (leaving aside gravity) obey the same equations of motion as light particles in a uniform waveguide. The relation: $E = H(p)$ becomes: $\hbar\omega = H(\hbar k)$, where $\hbar$ is a universal constant with the dimension of an action, ω is the (angular) frequency and k the wave number. In both cases, the product of the group velocity v and the phase velocity $\omega/k = E/p$ is unity. Thus the relation $H(p) = \sqrt{1 + p^2}$ postulated above may be understood as the one applicable to a wave propagating in a two dimensional space (say x, z), with boundary conditions on x . The corpuscle-wave analogy has been discussed in many works, see for example [24].

Gravity modeled by radiation pressure: Let us quote Feynman [1, p. 79]: “Many mechanisms for gravitation have been suggested. It is interesting to consider one of these. At first, one is quite excited and happy when he discovers it, but he soon finds that it is not correct. Suppose there were many particles moving in space at a very high speed in all directions and being only slightly absorbed when going through matter. When they are absorbed, they give an impulse to the earth.

However, since there are as many going one way as another, the impulse all balance. But when the sun is nearby, the particles coming toward the earth through the sun are partially absorbed, so fewer of them are coming from the sun than are coming from the other side. Therefore, the earth feels a net impulse toward the sun. What is wrong with that machinery: The earth would feel a resistance to motion [in free space], a phenomenon which is not observed". Note that if we hypothesize that the (zero mass, non self-interacting) particle speed is much larger than c (hypothesis whose compatibility with special relativity remains to be established) and replace absorption by some low-probability scattering, this gravity model would not contradict classical observations. In particular, if different corpuscles differ from one another only by different numbers of identical particles, the proportionality of w and m (constancy of g) would follow from the model.

The model presented below for a flat earth gravity is similar to the one described by Feynman, except that the particles that we consider (photons) are moving downward only and are totally reflected upon impinging on the corpuscle instead of being absorbed. The Feynman conclusion still applies: our model is accurate only when the corpuscle speed remains much smaller than the speed of light.

Consider an elastic collision between a corpuscle described by $H = H(p)$ and a particle described by $D = D(P)$, where H, D are the energies and p, P the momenta. Conservations of the total momentum and energy read:

$$\begin{aligned} p_1 + P_1 &= p_o + P_o && \text{conservation of momentum from } z \text{ invariance} \\ H(p_1) + D(P_1) &= H(p_o) + D(P_o) && \text{conservation of energy from } t \text{ invariance} \end{aligned} \quad (14)$$

Subscripts "o" refer to the particles before collision and subscripts "1" to particles after collision. When the particle is a photon we have, setting $c = 1$: $D(P) = |P|$, that is $D = -P$ when $P < 0$ (incident photons), and $D = P$ when $P > 0$ (reflected photons). Then, introducing the first equation in (14) into the second equation to eliminate P_1 that is not of interest to us, we obtain:

$$H(p_1) - p_1 = H(p_o) - p_o + 2P. \quad (15)$$

We have considered identical incident photons and set: $-P_o \equiv P > 0$. All the incident photons are assumed to have the same momentum. If P is small individual collisions do not affect much the corpuscle motion but a large number of collisions importantly modifies the corpuscle trajectory.

For a corpuscle of mass $m = 1$ we have: $H(p) = \sqrt{1 + p^2}$ as we have seen in the previous paragraph. If we introduce a parameter a defined by: $p \equiv \frac{1/a - a}{2}$, we have: $H(p) - p = a$. Then (15) reads: $a_1 = a_o + 2P$. Iterating for k collisions: $a_k = a_o + 2Pk$, $k = 0, 1, 2, \dots$. The corpuscle velocity $v = H'(p) = \frac{p}{\sqrt{1 + p^2}} = \frac{1 - a^2}{1 + a^2}$, that is: $v_k = \frac{1 - a_k^2}{1 + a_k^2}$, where v_o denote the initial velocity at $z = t = 0$ and v_1 denotes the velocity after the first collision. The velocity vanishes when $a_k = 1$, that is when: $k \approx \frac{1 - a_o}{2P}$.

We now suppose that the incident photons are spatially separated by a distance unity, in which case P represents the optical-wave incident power. If the corpuscle velocity is v , the time duration between two successive collisions is: $\Delta t = 1/(1 + v)$ from the definition of v , and: $\Delta t_k = \frac{1}{1 + v_k} = \frac{1 + a_k^2}{2}$. The altitude increment $\Delta z_k = v_k \Delta t_k = \frac{v_k}{1 + v_k} = \frac{1 - a_k^2}{2}$. Thus, Δt_k and Δz_k are simple functions of $a_k = a_o + 2kP$. The time t_n of occurrence of the n^{th} collision and its altitude z_n are

respectively:

$$\begin{aligned}
t_n &= \sum_{k=0}^n \Delta t_k = \sum_{k=0}^n \frac{1+a_k^2}{2} = \frac{n+1}{2} + s_n & z_n &= \sum_{k=0}^n \Delta z_k = \sum_{k=0}^n \frac{1-a_k^2}{2} = \frac{n+1}{2} - s_n \\
s_n &= \frac{1}{2} \sum_{k=0}^n (a_0 + 2Pk)^2 = \frac{a_o^2}{2} \sum_{k=0}^n 1 + 2a_o P \sum_{k=0}^n k + 2P^2 \sum_{k=0}^n k^2 \\
&= a_o^2 \left(\frac{n}{2} + \frac{1}{2} \right) + 2a_o P \left(\frac{n^2}{2} + \frac{n}{2} \right) + 2P^2 \left(\frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{6} \right)
\end{aligned} \tag{16}$$

For $P = 0$ (no radiation pressure) we have of course: $z_n/t_n = v_o$. For large n -values (small P -values):

$$Pt_n \approx x \left(\frac{1+a_o^2}{2} + a_o x + \frac{2x^2}{3} \right) \quad Pz_n \approx x \left(\frac{1-a_o^2}{2} - a_o x - \frac{2x^2}{3} \right) \tag{17}$$

In the approximate form given above only the product $x \equiv Pn$ is involved. We verify that for large x -values (large times) the corpuscle velocity tends to -1 since it is dragged down by the incident optical wave.

When the initial velocity $v_o = 1$ (ultra-relativistic corpuscle: $a_o = 0$), we easily find that the maximum altitude reached $z_m = 1/3P$. The return velocity, defined as the corpuscle velocity when $z = 0$ again, is $-1/2$. Since the corpuscle+light wave system is conservative, the energy lost by the corpuscle is delivered to the reflected optical wave, which we have not considered in the above discussion. When the initial velocity $v \ll 1$ we may set $a_o \equiv 1 - \delta$, $\delta \ll 1$. Then, to first order in δ we have: $v \approx p \approx \delta$, and we need not distinguish momentum and velocity (remember that $m = 1$). We also have from the preceding equations: $\Delta v \approx 2P$, $\Delta t \approx 1$ so that $Pt \approx Pn = x$. We are interested only in small x -values so that we may neglect the $2x^2/3$ term in (17). The acceleration $g = \Delta v / \Delta t = 2P$ being a constant we obtain, as from galilean mechanics: $z(t) = v_o t - g t^2 / 2 = \delta t - P t^2$. If we take into account the $2x^2/3$ term in (17) we find that the relative corpuscle energy loss over a period is nearly equal to $v_o \ll 1$ and is therefore negligible if we let c go to ∞ . However the $z(t)$ motion is not quite symmetrical in time.

From this paragraph, we conclude that the main text results apply to the case of gravity modelled by radiation pressure as long as the corpuscle speed is small compared with the speed of light. We have used the principle of invariance of the momentum and energy, but we have not postulated the law of dynamics: $dp/dt = w$, where w denotes the corpuscle weight.

Uniformly accelerated frame of reference: The Hamilton equations of motion in one space dimension (z) derive from a hamiltonian function of z and p (see Louis-Martinez [10], Eq(5) with $N = 1$):

$$H(z, p) = (mc^2 + m g z) \sqrt{1 + (p/mc)^2} \tag{18}$$

$$v(t) \equiv \frac{dz(t)}{dt} = \frac{\partial H(z, p)}{\partial p} = c \left(1 + \frac{z}{z_g} \right) \frac{\tilde{p}}{\sqrt{1 + \tilde{p}^2}} \quad \tilde{p} \equiv \frac{p}{mc} \quad z_g \equiv \frac{c^2}{g} \tag{19}$$

$$\frac{dp(t)}{dt} = - \frac{\partial H(z, p)}{\partial z} = -m g \sqrt{1 + \tilde{p}^2} \quad \rightarrow \quad \frac{d\tilde{p}(t)}{dt} = - \frac{g}{c} \sqrt{1 + \tilde{p}^2}. \tag{20}$$

We take the time origin such that: $p(0) = 0$. Note that the term $mc^2 + m g z_m \equiv mc^2(1 + z_m/z_g)$, where the maximum altitude z_m represents the corpuscle energy. The round-trip time is now a

function of $\zeta_m - \zeta$, where $\zeta = \ln(1 + z/z_g)$. The non-relativistic expressions are recovered when $z, ct \ll z_g$. An Hamiltonian function that would account for Quantum Gravity has been proposed by Chandra [25].

C The barometric law

We are considering an unity-area cylinder with vertical axis in uniform gravity resting on the ground: $z = 0$. We consider the motion along the vertical z -axis of a single corpuscle, denoted in general by $z = z(t; E)$, where t denotes time. The corpuscle energy is defined as: $E \equiv w z_m$ where w is the corpuscle weight and z_m the maximum altitude reached.

Consider a single period of corpuscle motion as shown in Fig. 1, without the dashed line. Let the corpuscle distance from the top of its trajectory be denoted by $Z \equiv z_m - z \geq 0$, at times t_1 and $t_2 \geq t_1$. We call: “round-trip time” $\tau(Z) \equiv t_2 - t_1$. Because in the present model gravity is static and uniform (that is, independent of time and altitude, respectively) this τ -function depends only on Z . As an example, for non-relativistic motion: $Z(t) = \frac{1}{2}gt^2$. The corpuscle weight $w \equiv mg$, where m denotes the corpuscle mass and g the gravitational acceleration. In that example $t_{1,2} = \pm\sqrt{2Z/g}$ and thus $\tau(Z) = 2\sqrt{2Z/g} \propto \sqrt{Z/w}$, m being a constant. For a unity weight we may take simply: $\tau(Z) = \sqrt{Z}$. The period of motion of a corpuscle bouncing off the ground ($z = 0$) without any loss of energy (rigid walls and negligible gas friction) and having energy z_m , is according to the above definitions: $\tau(z_m)$. The time spent by the corpuscle above some z -level is obviously zero if $z_m < z$. If $z_m \geq z$ this time is: $\tau(z_m - z)$ as suggested on Fig. 1.

We now suppose that the ground on which the corpuscle is bouncing off is in contact with a heat bath. We are implicitly considering the so-called “canonical ensemble”: the ground plate does not allow corpuscles to go through, but it moves slightly so that a transfer of energy between the bath and the corpuscle is allowed. This means that the ground level ($z=0$) is not perfectly static as assumed above, but instead is quivering. Concretely, the ground level could be moving up and down according to some zero-mean fluctuation of small amplitude so that, upon impacting on the ground, the corpuscle may loose energy (when the ground level is moving downward), or gain energy (when the ground level is moving upward). We will not need the explicit form of this fluctuation. We only assume that the amplitude of that fluctuation is so small that the corpuscle energy does not vary significantly over many periods. Yet it may evolve slowly. The energy distribution $\omega(z_m)$ refers to averages over arbitrarily long times, and is presently unknown; it will be determined by enforcing the condition that the law of interest (presently the barometric law) does not depend on the laws of physics (except the law of conservation of energy, which follows from the time-translation invariance of the system). On the average, the fraction of time $\mathcal{B}$ during which the corpuscle is above some z -level is:

$$\mathcal{B}(z) = \frac{\int_z^\infty dz_m \omega(z_m) \tau(z_m - z)}{\int_0^\infty dz_m \omega(z_m) \tau(z_m)}. \quad (21)$$

We now argue that $\omega(z_m)$ *must be*: $\exp(-w z_m/\theta)$, where θ is a constant having the dimension of an energy, as is required by the fact that the argument of an exponential is dimensionless. First,

let us introduce this distribution law in (21). We obtain:

$$\begin{aligned}
\mathcal{B}(z) &= \frac{\int_z^\infty dz_m \exp(-w z_m/\theta) \tau(z_m - z)}{\int_0^\infty dz_m \exp(-w z_m/\theta) \tau(z_m)} \\
&= \frac{\exp(-w z/\theta) \int_z^\infty dz_m \exp(-w (z_m - z)/\theta) \tau(z_m - z)}{\int_0^\infty dz_m \exp(-w z_m/\theta) \tau(z_m)} \\
&= \frac{\exp(-w z/\theta) \int_0^\infty dz'_m \exp(-w z'_m/\theta) \tau(z'_m)}{\int_0^\infty dz_m \exp(-w z_m/\theta) \tau(z_m)} \\
&= \exp(-w z/\theta).
\end{aligned} \tag{22}$$

On the third line, we have used as an integration variable $z'_m \equiv z_m - z$ instead of z_m . The last line follows from the fact that z_m, z'_m are dummy variables so that we may replace z'_m by z_m . Note that, even though we have introduced integral signs, no integration has been performed. We have only used the fact that: $\int_a^\infty dx f(x - a) = \int_0^\infty dy f(y)$, which follows from the definition of an integral as a sum of terms and the associativity of real-number addition.

The distribution $\omega(z_m)$ introduced above is the only one that leads to a result (last line of (22)) independent of the particular form of the $\tau(Z)$ -function, and therefore of the law of motion. The only function of $u \equiv w z_m/\theta$ that may cause the integral in the numerator to go from 0 to infinity and cancel out with the integral in the denominator is: $\exp(-u)$, the argument being defined only to within a constant factor. Indeed, the exponential $f(x) = \exp(x)$ is the only function such that $f(a + b) = f(a)f(b)$ for any a, b (of course, the function 2^x that would have been more easily comprehended by the ancient greeks, is equally acceptable). In order to obtain the distribution $\omega(z_m)$, one would need to know the function $\tau(Z)$. But, remarkably, this distribution is not needed.

The gas density, defined as the probability that the corpuscle be located between z and $z + dz$, divided by dz , is: $\rho(z) = -d\mathcal{B}(z)/dz = (w/\theta) \exp(-w z/\theta)$. This is the barometric law. Since w and θ are constants the density decays exponentially as a function of altitude z . In the earth atmosphere the density of di-atomic oxygen decays faster than the density of di-atomic nitrogen because the weight of an oxygen atom exceeds that of a nitrogen atom in the ratio $\approx 16/14$.

Two weight values: We now suppose that the weight is w_1 for $0 < z \leq h$ and w_2 for $z \geq h$. We evaluate the ratio $\mathcal{B}(h)$ of the average time spent by the corpuscle above the h altitude and the average period. Let us denote $\tau_i(Z)$ the round-trip time corresponding to a distance Z from the top of the trajectory when the weight is a constant w_i with $\tau_i(0) = 0$. These functions are different for different i . But continuity of the corpuscle motion entails that: $\tau_2(Z) = \frac{w_1}{w_2} \tau_1(\frac{w_2}{w_1} Z)$, as one can see by taking the derivative with respect of Z . For non-relativistic motion $\tau_i(Z) = C \sqrt{Z/w_i}$, where C is a constant, and the above relation is readily verified. As in the main text E denote the corpuscle energy and we consider a motion period. When $E \leq w_1 h$ the corpuscle stays below the h -plane and therefore the duration above h is equal to zero. When $E > w_1 h$ the time spent by the corpuscle above the h -plane is $\tau_2(\frac{E - w_1 h}{w_2})$ because the corpuscle weight is then w_2 and its energy with respect to the h -plane is $E - w_1 h$. Thus the average time spent above the h -plane is:

$$\text{average time above } h = \int_{w_1 h}^\infty dE \omega(E) \tau_2\left(\frac{E - w_1 h}{w_2}\right) = \int_{w_1 h}^\infty dE \omega(E) \frac{w_1}{w_2} \tau_1\left(\frac{E}{w_1} - h\right) \tag{23}$$

where we have used the relation between $\tau_1(Z)$ and $\tau_2(Z)$ given above.

The evaluation of the average period is more complicated. We find:

$$\text{average period} = \int_0^{w_1 h} dE \omega(E) \tau_1\left(\frac{E}{w_1}\right) + \int_{w_1 h}^{\infty} dE \omega(E) \left(\frac{w_1}{w_2} \tau_1\left(\frac{E}{w_1} - h\right) + \tau_1\left(\frac{E}{w_1}\right) - \tau_1\left(\frac{E}{w_1} - h\right) \right). \quad (24)$$

When $E \leq w_1 h$ (first integral in the above expression) the corpuscle does not reach the $z = h$ altitude, and thus is all the time below h . Accordingly, since $Z = E/w_1$, the time spent below h is $\tau_1(\frac{E}{w_1})$. Consider now the second integral, $E > w_1 h$. The first term is the time spent above h given in the preceding equation (23). To evaluate the time spent below h , consider the corpuscle motion as if the weight were w_1 everywhere ($0 < z < \infty$). Then we would have $z_m = E/w_1$ and the time spent below $z = h$ is: $\tau_1(z_m) - \tau_1(z_m - h)$. The corpuscle motion below h must be split time-wise to meet the condition of continuity of $z(t)$ at $z = h$. However, this splitting does not affect the difference: $\tau_1(z_m) - \tau_1(z_m - h)$.

It follows from (23) and (24) that:

$$\begin{aligned} \mathcal{B}(h) &= \frac{\int_{w_1 h}^{\infty} dE \omega(E) \frac{w_1}{w_2} \tau_1\left(\frac{E}{w_1} - h\right)}{\int_0^{w_1 h} dE \omega(E) \tau_1\left(\frac{E}{w_1}\right) + \int_{w_1 h}^{\infty} dE \omega(E) \left(\frac{w_1}{w_2} \tau_1\left(\frac{E}{w_1} - h\right) + \tau_1\left(\frac{E}{w_1}\right) - \tau_1\left(\frac{E}{w_1} - h\right) \right)} \\ &= \frac{\frac{w_1}{w_2} \int_{w_1 h}^{\infty} dE \omega(E) \tau_1\left(\frac{E}{w_1} - h\right)}{\int_0^{\infty} dE \omega(E) \tau_1\left(\frac{E}{w_1}\right) + \left(\frac{w_1}{w_2} - 1\right) \int_{w_1 h}^{\infty} dE \omega(E) \tau_1\left(\frac{E}{w_1} - h\right)} \end{aligned} \quad (25)$$

The only way to remove the τ -functions, according to the principle of simplicity postulated in the main text is to choose: $\omega(E) = \exp(-E/\theta)$, where θ has the dimension of an energy. We obtain from the above expression:

$$\mathcal{B}(h) = \frac{1}{1 + \frac{w_2}{w_1} \left(\exp\left(\frac{w_1 h}{\theta}\right) - 1 \right)}. \quad (26)$$

We have therefore obtained an expression applicable to a non-linear potential. We recover the previous result in (22) when $w_1 = w_2 \equiv w$. Of course, when $h = 0$, $\mathcal{B} = 1$: the corpuscle is always at some altitude.

The expression in (26) is usually obtained by postulating a gas density: $\rho(z) \propto \exp(-\phi(z)/\theta)$, where $\phi(z)$ is the potential at z : $\phi(z) = w_1 z, z \leq h$ and $\phi(z) = w_1 h + w_2(z - h), z > h$. $\mathcal{B}$ is then the ratio of the integral of $\rho(z)$ from h to ∞ and its integral from 0 to ∞ . In our formulation no integration is needed.

Another example: The weight is $1/\Delta_1$ for $0 < z < \Delta_1 \equiv z_1$, and $1/\Delta_2$ for $\Delta_1 < z < \Delta_1 + \Delta_2 \equiv z_2$. According to the traditional method, the probability that the corpuscle be in the upper section is: $s_2/(s_1 + s_2)$ where:

$$s_1 \equiv \int_0^{z_1} dz \exp\left(-\frac{z/z_1}{\theta}\right) = \int_0^{\Delta_1} dz \exp\left(-\frac{z/\Delta_1}{\theta}\right) = \theta(1 - x)\Delta_1 \quad x \equiv \exp(-1/\theta) \quad (27)$$

For $z_1 < z < z_2$ we have: $\phi(z) = 1 + (z - z_1)/(z_2 - z_1)$. Setting $\zeta \equiv z - z_1$:

$$s_2 \equiv \int_{z_1}^{z_2} dz \exp\left(-\frac{1 + (z - z_1)/(z_2 - z_1)}{\theta}\right) = \int_0^{\Delta_2} d\zeta \exp\left(-\frac{1 + \zeta/\Delta_2}{\theta}\right) = \theta(1 - x)\Delta_2 x. \quad (28)$$

Thus:

$$\mathcal{B}(h) = \frac{s_2}{s_1 + s_2} = \frac{1}{1 + \frac{\Delta_1}{\Delta_2} \exp(\frac{1}{\theta})}. \quad (29)$$

Let us now consider the method presented in the main text. For $0 < E < 1$ the corpuscle does not reach the intermediate plane, for $1 < E < 2$ it crosses the intermediate plane but does not reach the upper plane, and for $E > 2$ it is reflected by the upper plane. We need evaluate the average times when $z = z_1$ and the average time when $z = 0$.

	$z = 0$	$z = z_1$
$E < 1$	$\Delta_1 \tau(E)$	0
$1 < E < 2$	$\Delta_2 \tau(E - 1) + \Delta_1 (\tau(E) - \tau(E - 1))$	$\Delta_2 \tau(E - 1)$
$2 < E$	$\Delta_2 (\tau(E - 1) - \tau(E - 2)) + \Delta_1 (\tau(E) - \tau(E - 1))$	$\Delta_2 \tau(E - 1) - \Delta_2 \tau(E - 2))$

(30)

The integrated duration at z_1 is:

$$T_1 = \int_1^2 dE \omega(E) \Delta_2 \tau(E - 1) + \int_2^\infty dE \omega(E) (\Delta_2 \tau(E - 1) - \Delta_2 \tau(E - 2)) \quad (31)$$

If $\omega(E) = \exp(-E/\theta)$ we obtain through transformation similar to the ones made earlier:

$$T_1 = \Delta_2 x (1 - x) \int_0^\infty dE \exp(-E/\theta) \tau(E) \quad (32)$$

Similarly:

$$\begin{aligned} T_0 &= \int_0^1 dE \omega(E) \Delta_1 \tau(E) \\ &+ \int_1^2 dE \omega(E) (\Delta_2 - \Delta_1) \tau(E - 1) + \int_1^2 dE \omega(E) \Delta_1 \tau(E) \\ &+ \int_2^\infty dE \omega(E) (\Delta_1 \tau(E) + \Delta_2 \tau(E - 1) - \Delta_2 \tau(E - 2)) \\ &= (\Delta_1 + (\Delta_2 - \Delta_1)x - \Delta_2 x^2) \int_0^\infty dE \exp(-E/\theta) \tau(E) \end{aligned} \quad (33)$$

Thus:

$$\mathcal{B} = \frac{T_1}{T_0} = \frac{\Delta_2 x}{\Delta_1 + \Delta_2 x}, \quad (34)$$

which coincides with the result in (29) obtained by the traditional method.

D A report to the “American Journal of Physics”

The present paper was submitted to the American Journal of Physics. The editor of that journal received the following report: “The authors propose to establish the barometric formula in particular, or the ideal gas law in general, straight from atomistic principles (à la Democrite), independent

of any particular laws of motion. True, in most text-books this follows as an afterthought, after the derivation of Boltzmann, which may be overkill. (A minor exception to this, which I always found cute, is the derivation of the Maxwell distribution in O’Hanian’s book on modern physics, which is based only on the isotropy of space but relies on a purely classical notion of velocity and I don’t know whether it can be generalized.) Assuming only that the time spent in a region is related to the potential against which a particle would have to climb, which follows from the energy-work theorem, an exponential distribution is obtained as a function of height, independent of whether the motion is relativistic or not. The presentation is clear and insightful, which will greatly benefit the reader, and inspire similar thought on other taken-for-granted assumptions. I can not judge, though, how original these ideas are. I am almost certain that there are many insiders for whom this would be obvious, but I am not sure whether anybody would have written it down in such a pedagogical and thoughtful manner.”

Note 1: The second edition of the book by O’Hanian’s cited in this report does not seem to involve concepts similar to the ones presented in our paper.

Note 2: A second AJP referee asserted that the present paper results are erroneous and that, accordingly, our paper should be rejected. We have explained in a previous “arxiv” version why this second referee assertion is unfounded. As emphasized in the paper present version (identical in basic content to the previous versions), our results are in exact agreement with all papers and books (e.g., Greiner, cited) known to us dealing with the special relativity ideal-gas solution in the presence (or not) of gravity.

References

- [1] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman lectures in physics*. Reading: Addison-Wesley, 1977, sixth printing.
- [2] According to *general relativity*, thermal energy has weight. But this (so-called Tolman) effect that entails a temperature variation on the order of 10^{-12} on earth is entirely negligible.
- [3] J. Arnaud, L. Chusseau, and F. Philippe, “A simple model for Carnot heat engines,” *Am. J. Phys.*, vol. 78, no. 1, pp. 106–110, 2010. [Online]. Available: <http://arxiv.org/abs/0812.2757>
- [4] J. Arnaud, L. Chusseau, and F. Philippe, “Démocrate et la puissance motrice du feu (Democritus and the motive power of fire),” 2011, 79 pages. [Online]. Available: <http://arxiv.org/abs/1104.0836>
- [5] Concretely, the corpuscle may be a cold neutron submitted to the earth gravity and reflected by a diamond at some non-zero temperature [26].
- [6] F. L. Roman, J. A. White, and S. Velasco, “Microcanonical single-particle distributions for an ideal gas in a gravitational field,” *European Journal of Physics*, vol. 16, p. 83, 1995.
- [7] H. C. Callen, *Thermodynamics and a introduction to thermo-statistics*, 2nd ed. New-York: Wiley and Sons, 1985.
- [8] P. T. Landsberg, J. Dunning-Davies, and D. Pollard, “Entropy of a column of gas under gravitation,” *Am. J. Phys.*, vol. 62, no. 8, pp. 712–717, 1994.

- [9] R. Leinonen, E. Raesaenen, M. Asikainen, and P. E. Hirvonen, “Students’ pre-knowledge as a guideline in the teaching of introductory thermal physics at university,” *European Journal of Physics*, vol. 30, pp. 593–604, 2009.
- [10] D. J. Louis-Martinez, “Classical relativistic ideal gas in thermodynamic equilibrium in a uniformly accelerated reference frame,” arXiv 1012.3063, Dec. 2010. [Online]. Available: <http://arxiv.org/abs/1012.3063>
- [11] Note that: $2 \int_0^\infty dx x \exp(-x) \sqrt{x} = 3 \int_0^\infty dx \exp(-x) \sqrt{x}$.
- [12] W. Greiner, L. Neise, and H. Stöcker, *Thermodynamics and Statistical Mechanics*. New-York: Springer-Verlag, 1997.
- [13] D. Levermore, “Early theories of gases,” <http://www2.math.umd.edu/~lvrmr/History/EarlyTheories.html>.
- [14] E. Hecht, *Physique*. de Boeck University, 2003, see § 14.8.
- [15] J. Waterston, “On the physics of media that are composed of free and perfectly elastic molecules in a state of motion,” *Philosophical Transactions of the Royal Society of London A*, vol. 183, pp. 1—79, 1892, this article, submitted to that journal in 1843, was initially rejected.
- [16] F. P. Miller, A. F. Vandome, and J. McBrewster, Eds., *Ideal gas law*. Mauritius: Alphascript Publishing, 2009, this booklet reproduces Wikipedia articles relating to the ideal-gas law. Numerous references to Physics textbooks are listed there. See also: http://en.wikipedia.org/wiki/Kinetic_theory.
- [17] We quote: Gases consist in corpuscles having non-zero mass. The corpuscles are quickly moving. They are perfectly spherical and elastic. The average kinetic energy depends only on the system temperature. Relativistic effects are negligible. Motion laws are time reversible. The number of molecules is so large that a statistical treatment is appropriate.
- [18] J. D. Norton, “Atoms, entropy, quanta: Einstein’s miraculous argument of 1905,” *Studies in History and Philosophy of Modern Physics*, vol. 37, pp. 71–100, 2006. [Online]. Available: <http://www.pitt.edu/~jd/papers/miraculous.pdf>
- [19] J. Arnaud, L. Chusseau, and F. Philippe, “On the ideal gas law,” arXiv 1105.2643 v2, May 2011. [Online]. Available: <http://arxiv.org/abs/1105.2643>
- [20] The time-independent hamiltonian form is given here for the readers convenience: If $H(z, p)$ denotes the hamiltonian function of coordinate z and momentum p , we have:

$$v(t) \equiv \frac{dz(t)}{dt} = \frac{\partial H(z, p)}{\partial p} \quad \frac{dp(t)}{dt} = -\frac{\partial H(z, p)}{\partial z}. \quad (35)$$

Because the above hamiltonian function does not depend explicitly on t , the energy of a corpuscle is the constant: $E = H(z, p)$ for some initial z, p values. One readily verifies that the total derivative: $dE/dt = 0$. The main-text expression corresponds to: $H(z, p) = H(p) + w z$.

- [21] Indeed, the equation $z(t) - z_o = -\frac{H(-w(t-t_o))}{w}$ shows that $\tilde{t} \equiv wt$ is a universal function of $\tilde{z} \equiv wz$, that is, this relation does not depend on w . This implies that if $w_1 z_1 = w_2 z_2$ then $w_1 t_1 = w_2 t_2$. Thus $t_2 = (w_1/w_2)t_1(z_1) = (w_1/w_2)t_1((w_2/w_1)z_2)$. To obtain Eq. 12, set $z_2 \equiv Z$, and $t_1 - > \tau_1$, $t_2 - > \tau_2$.

- [22] L. D. Landau and E. M. Lifshitz, *Statistical Physics*. Oxford: Pergamon Press, 1980, vol. 5, Part I.
- [23] The traditional methods of calculating $\langle F \rangle$ and U in special relativity but in the absence of gravity consist instead in calculating the average values of $p v$ and $H(p)$, respectively, using as a weight the Boltzmann factor: $\exp(-H(p)/\theta)$.
- [24] J. A. Arnaud, *Beam and Fiber Optics*. Academic Press, 1976.
- [25] N. Chandra and S. Chatterjee, “Thermodynamics of ideal gas in doubly special relativity,” arXiv 1108.0896, Aug. 2011. [Online]. Available: <http://arxiv.org/abs/1108.0896>
- [26] H. Kaneko, A. Tohsaki, and A. Hosaka, “Thermodynamics of quantum ultra-cold neutron gas under gravity of the earth,” arXiv:1205.0378v1 [quant-ph], May 2012. [Online]. Available: <http://arxiv.org/abs/1205.0378>